



Published in final edited form as:

*Am J Geriatr Psychiatry*. 2015 May ; 23(5): 466–476. doi:10.1016/j.jagp.2014.10.005.

## Anxiety symptoms in amnesic mild cognitive impairment are associated with medial temporal atrophy and predict conversion to Alzheimer's disease

Linda Mah, MD, MHS<sup>1,2</sup>, Malcolm A. Binns, PhD<sup>1,3</sup>, David C. Steffens, MD, MHSc<sup>4</sup>, and the Alzheimer's Disease Neuroimaging Initiative \*

<sup>1</sup>Rotman Research Institute, Baycrest, Toronto, Canada

<sup>2</sup>Division of Geriatric Psychiatry, University of Toronto, Canada

<sup>3</sup>Dalla Lana School of Public Health, University of Toronto, Canada

<sup>4</sup>Department of Psychiatry, University of Connecticut Health Center, Farmington, CT, USA

### Abstract

**Background**—The purpose of the current study was to test the hypothesis that anxiety in amnesic mild cognitive impairment (aMCI) increases rates of conversion to Alzheimer's disease (AD), and to identify potential neural mechanisms underlying such an association.

**Methods**—376 participants with aMCI from the Alzheimer's Disease Neuroimaging Initiative (ADNI) were studied over a median period of 36 months. A Cox proportional-hazards model was used to assess the association between anxiety severity ratings on the Neuropsychiatric Inventory Questionnaire and AD risk. Other variables in the model were depression, memory loss, and MRI-derived AD-related regions-of-interest (ROI), including hippocampal (HC), amygdalar (AMYG), entorhinal cortical (EC) volumes and EC thickness. In addition, a linear regression model was used to determine the effect of anxiety in aMCI on rates of atrophy within ROIs.

**Results**—Anxiety severity increased rate of aMCI conversion to AD, after controlling for depression and cognitive decline. The association between anxiety and AD remained significant

© 2014 American Association for Geriatric Psychiatry. All rights reserved.

Corresponding Author: Linda Mah, MD, MHS, FRCPC Rotman Research Institute, Baycrest Division of Geriatric Psychiatry, University of Toronto 3560 Bathurst Street Toronto, Ontario M6A 2E1 Phone: (416) 785-2500 ext. 3365 Fax: (416) 785-4230 [lmah@research.baycrest.org](mailto:lmah@research.baycrest.org).

\*Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database ([adni.loni.ucla.edu](http://adni.loni.ucla.edu)). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: [http://adni.loni.ucla.edu/wp-content/uploads/how\\_to\\_apply/ADNI\\_Acknowledgement\\_List.pdf](http://adni.loni.ucla.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf)

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

#### AUTHOR CONTRIBUTIONS

All authors contributed to drafting of the manuscript, revising it critically for important intellectual content, and approved the final manuscript. LM, MB, DS: conception and design of the study, interpretation of results, critical review. MB: Statistical analyses.

#### CONFLICT OF INTEREST DISCLOSURES

None of the authors reported potential conflicts of interest.

even with inclusion of ROI baseline values or atrophy rates as explanatory variables. Further, anxiety status predicted greater rates of decrease in EC volume. An association between anxiety and EC-thickness missed significance.

**Conclusions**—Anxiety symptoms in aMCI predict conversion to AD, over and beyond the effects of depression, memory loss, or atrophy within AD neuroimaging biomarkers. These findings, together with the greater EC atrophy rate predicted by anxiety, are compatible with the hypothesis that anxiety is not a prodromal non-cognitive feature of AD, but may accelerate decline towards AD through direct or indirect effects on EC.

### Keywords

Alzheimer's disease; anxiety; depression; mild cognitive impairment; neuropsychiatric symptoms; amygdala; entorhinal cortex; MRI biomarker

## INTRODUCTION

Amnesic Mild Cognitive Impairment (aMCI) is characterized by memory impairment with preservation of functional independence and is considered a transitional stage between normal aging and Alzheimer's disease (AD)(1-3). However, rates of conversion to AD are highly variable. While 10-15% of aMCI convert to AD within a year, others remain stable or improve in memory performance(3, 4). The majority of models incorporating cerebrospinal fluid, neuroimaging, and neuropsychological biomarkers have inadequate predictive value in identifying aMCI at high risk for disease progression(5-8). These findings suggest that other variables contribute to AD risk.

Neuropsychiatric symptoms, such as depression and anxiety, are frequent in aMCI(9-13). Much research has focused on the link between depression and AD, with converging evidence of a 1.5-2 times increased risk of conversion in individuals with aMCI and concurrent depression(14). Whether anxiety is a risk factor for AD is unclear. Past or current anxiety in aMCI may augment the risk of AD conversion by as much as 2.5 fold(15-18). Conversely, anxiety may be protective against AD(19-21) or have no impact(22-25). Although methodological differences such as shorter follow-up(22-25) and clinical versus community samples(19-21) may account for the discrepant findings, an important consideration is the potential impact of concurrent depression, given the well-established comorbidity between anxiety and depression in the general population(26, 27). Notably, studies which included depression as a covariate reported an inverse or no association between anxiety and AD(19-21, 25). One of the first studies to demonstrate a nearly two-fold increased AD risk in individuals with aMCI and anxiety reported that anxiety at baseline remained a significant predictor when depression was included as a covariate, although the results were not presented(17). Other studies which demonstrated an association between aMCI with anxiety and conversion excluded depression as a covariate in longitudinal models of AD risk(15, 16, 18). Thus, while evidence from large observational studies supports an association between anxiety in aMCI and AD risk, the degree to which depression contributes towards, or confounds the association is unclear.

An association between anxiety and AD risk is suggested by studies which have demonstrated links between anxiety or stress and AD-related pathology. Anxiety in individuals with aMCI was associated with abnormal concentrations of A $\beta$ 42 and t-tau in cerebrospinal fluid, while conversely, depression and apathy were not(28). In animal models of AD, stress-level glucocorticosteroid administration resulted in increases in amyloid formation and tau accumulation(29). The well-established relationships amongst stress, cortisol, and hippocampal atrophy in humans(30-32) suggest that anxiety may increase AD risk indirectly through cortisol-mediated hippocampal neurotoxicity. Elevated cortisol levels have been found in older adults with clinically significant anxiety(33). Chronic stress may also contribute to AD risk through other pathways related to increased allostatic load, such as effects on the cardiovascular system and plasticity changes to brain structures(34).

The purpose of the current study was to test the hypothesis that anxiety symptoms in individuals with aMCI increase risk of AD, independent of depressive symptoms. We also sought to identify potential neural mechanisms underlying such an association by determining the relationships amongst anxiety symptoms in aMCI, AD-related brain regions, and conversion to AD. Medial temporal lobe structures, which together play an integral role in memory formation, and are robust predictors of progression to AD(8), were selected as regions-of-interest. These were hippocampal (HC-vol), amygdala (AMYG-vol), and entorhinal cortex volumes (EC-vol) as well as cortical thickness measures (EC-thickness). We additionally included EC surface area measures (EC-surface) based on neuropathological studies of early AD(35). We hypothesized that anxiety symptoms in aMCI (aMCI anxiety+) would be associated with greater rates of atrophy within ROIs relative to non-anxious aMCI (aMCI anxiety-).

## METHODS AND MATERIALS

### Subjects

Data used in the preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative(ADNI) database(36). ADNI was launched in 2003 by the NIA, National Institute of Biomedical Imaging and Bioengineering, the FDA, private pharmaceutical companies, and nonprofit organizations as a \$60M, five-year public-private partnership. The primary goal of ADNI has been to test whether serial magnetic resonance imaging(MRI), positron emission tomography, other biological markers, and clinical and neuropsychological assessment can be combined to measure the progression of MCI and early AD.

The Principal Investigator of this initiative is Michael W. Weiner, MD, VA Medical Center, UCSF. ADNI is the result of efforts of many co-investigators from a broad range of academic institutions and private corporations. Subjects have been recruited from over 50 sites across the USA and Canada. The initial goal of ADNI was to recruit 800 subjects. ADNI has been followed by ADNI-GO and ADNI-2. Subjects originally recruited for ADNI-1 and ADNI-GO had the option to be followed in ADNI-2(8). For up-to-date information, see [www.adni-info.org](http://www.adni-info.org).

Diagnoses were established by site investigators at screening and baseline visits using quantitative and qualitative global measures. General inclusion criteria were age 55–91 years for MCI and AD, 60–90 years for NC, availability of family/caregiver to provide collateral information, modified Hachinski ischemia score  $\leq 4$ , stability of permitted medications, no significant neurological conditions, and good general health. Excluded were individuals with active mood disorders such as major depressive or bipolar disorder, or those with Geriatric Depression Scale score  $> 5$  (37). Criteria for single- or multiple-domain aMCI (1) included memory complaint by subject or family/caregiver, objective memory loss (performance 1.5 standard deviations below education-adjusted cutoff scores on Wechsler Memory Scale-Revised (WMS-R) Logical Memory-IIa), no significant impairment in other cognitive domains, MMSE score  $\geq 24$  (38), Global Clinical Dementia Rating (CDR) score = 0.5 (39), CDR memory score  $\geq .5$ , preserved activities of daily living, and absence of dementia. Detailed inclusion and exclusion criteria are available in the ADNI protocol (36).

Clinical, neuropsychological and neuroimaging assessments were collected at baseline and at 6 month intervals for 2 years, then annually. Data were downloaded from the ADNI public database on or before February 09, 2014 and reflect the status of the database at that point. The ADNI study was approved by the local institutional review board of each participating site. Prior to performance of any study procedures, informed consent was obtained from all subjects and study partners.

The current study included 376 aMCI participants from ADNI-1 and was approved by Baycrest's REB.

### Assessment of exposure of interest

Anxiety in ADNI-1 was assessed using the Neuropsychiatric Inventory (NPI-Q) (40), a caregiver-based report of neuropsychiatric symptoms associated with dementia which have occurred over the last month. The NPI-Q uses a two-part screening question for anxiety, probing whether the patient becomes upset when separated from caregiver or whether patient has physiological symptoms of anxiety (e.g., shortness of breath, feeling tense). Endorsement of either is recorded as a positive response for anxiety. Severity of anxiety is then rated (1=mild, 2=moderate, 3=severe). NPI-Q responses were available for baseline and follow-up visits (every 6 months until 24 months, then annually). As a summary measure of exposure to anxiety during the observation period, we used the highest reported anxiety severity rating (NPI-Anxiety<sub>max</sub>) recorded during the period of risk, defined as baseline through to the study visit immediately prior to AD diagnosis, or to the second-to-last study visit in stable aMCI participants. Negative responses on NPI-Q anxiety were coded as 0 for severity.

**Other potential predictors of AD risk**—Baseline age, sex, education, Hachinski ischemia score (41), and the following depression, cognitive, and AD neuroimaging biomarkers were included as explanatory variables of AD risk.

**Depression**—Assessments included the Geriatric Depression Scale (GDS), administered at baseline and then yearly, and the NPI-Q. We used the highest GDS score (GDS<sub>max</sub>) and

NPI-Q depression severity rating (NPI-Dep<sub>max</sub>) recorded during the period of risk as measures of exposure to depression.

**Cognition**—Baseline Mini-Mental Status Examination (MMSE)(38) and neuropsychological measures of memory and executive function were included. ADNI Memory (ADNI-MEM) and Executive Function (ADNI-EF) are composite scores created to address site-to-site variability in neuropsychological test versions administered(42). ADNI-MEM is based on the Rey Auditory Verbal Learning Test (RAVLT)(43), AD Assessment Scale-Cognitive Subscale (ADAS-Cog)(44), MMSE 3-item recall, and WMS-R Logical Memory-I(45). ADNI-EF is based on Category Fluency (animals, vegetables), Trails B, Digit span backwards, WAIS-R Digit Symbol Substitution, and five Clock Drawing items. Each metric was constructed with a mean of 0 and standard deviation of 1 based on the pool of 800 participants who completed baseline neuropsychological assessment in ADNI-1. ADNI-MEM and ADNI-EF have been validated as predictors of AD risk and were associated with AD structural neuroimaging biomarkers(42, 46). ADNI-EF and ADNI-MEM scores at baseline and ADNI-MEM scores at each follow-up during the period of risk were incorporated into analyses.

**Structural neuroimaging ROIs**—Magnetic resonance imaging (MRI)-derived measures of total intracranial volume (ICV), HC-vol, AMYG-vol, EC-vol, EC-thickness, and EC-surface at baseline and at each study visit (every six months until 24 months, then annually) were extracted from the dataset UCSF-Longitudinal FreeSurfer (FreeSurfer Version 4.4). The dataset includes MRI measures of cortical volume and thickness, standard deviation of cortical thickness, and surface area for 34 ROIs. Participants were scanned on 1.5-Tesla scanners at 56 different sites using a standardized ADNI protocol for high-resolution MPRAGE structural MRI. Images were preprocessed using gradient warping, scaling, B1 correction and N3 inhomogeneity correction. Original scans and pre-processed images are available at (<http://www.loni.ucla.edu/ADNI/Data/>)(36). Cortical reconstruction and volumetric segmentation of scans was performed using FreeSurfer(<http://surfer.nmr.mgh.harvard.edu/>). Because MRI scans were acquired at multiple time-points, scans were run through FreeSurfer using longitudinal processing, which involves creating a within-subject template space and average image unbiased toward the chronological scan order using robust, inverse consistent registration(47). Several processing steps, including skull stripping, Talairach transforms, atlas registration as well as spherical surface maps and parcellations were then initialized with common information from the within-subject template to optimize statistical power and reliability(48). Further details on longitudinal processing are available here(48).

## Statistical Analyses

A Cox proportional-hazards regression model was used to evaluate the effects of anxiety and other factors on time to conversion to AD. Factors were NPI-Anxiety<sub>max</sub>, GDS<sub>max</sub>, NPI-Dep<sub>max</sub>, sex, education, baseline age, Hachinski ischemia score, ADNI-MEM, ADNI-EF, and ROIs. This model yields a hazard ratio (HR) which gives the ratio of the probability of converting to AD at a given time per 1 unit of the predictor (anxiety severity rating), relative to the probability of converting to AD for aMCI with no anxiety. For multivariable

regression models, we report the HR for the predictor conditional on other variables in the model ( $HR_p$ ). To account for variability in anxiety and depression severity across the observation period, NPI-Anxiety<sub>max</sub>, NPI-Dep<sub>max</sub>, and GDS<sub>max</sub> were entered as time-dependent variables (TDV)(49).

To appreciate the unconditional contribution of individual factors on AD risk, we fit bivariate Cox models with each factor as the independent variable. We then fit a multivariable Cox model using NPI-Anxiety<sub>max</sub> and all demographic and clinical factors to determine the association between anxiety and AD, conditional on other predictors of AD risk. We used backward elimination to sequentially remove variables with  $p > .10$ . This final model was confirmed by re-introducing each eliminated variable into the model to ensure  $p > .10$  and to determine whether the regression coefficient estimate for anxiety was substantively altered ( $\geq 20\%$ ) by inclusion of the variable.

To evaluate the extent to which anxiety was predictive of AD over and beyond extent of memory decline, we challenged the final model by including a measure of change in memory ( $\Delta$ ADNI-MEM) and determined its impact on the size of the regression coefficient estimate for anxiety.  $\Delta$ ADNI-MEM was calculated as the difference between ADNI-MEM at baseline and at each follow-up visit during the period of risk and then entered into the model as a TDV.

### Structural neuroimaging predictors of AD risk

MRI data were available for a subset of the sample ( $n=332$ ). To determine whether anxiety was predictive of AD conversion over and beyond the contribution of structural neuroimaging biomarkers of AD, we refit the Cox regression model to the subsample using the previously-described procedure. Baseline ICV was added to the model to account for variability in total brain volume across participants, then baseline values for each ROI were entered individually into the model to determine their impact on the regression coefficient estimate for anxiety.

The same procedure was followed in evaluating the impact of change in each ROI ( $\Delta$ ROI) over time in the model.  $\Delta$ ROI was calculated as the difference from baseline to each scan obtained during the period of risk.  $\Delta$ ICV was included in these models.

### Anxiety as a predictor of rate of atrophy within ROIs

To explore potential neural mechanisms underlying an association between anxiety and AD, we evaluated whether aMCI anxiety+ versus aMCI anxiety- status was predictive of rate of atrophy ( $R_{atrophy}$ ) within ROIs over the entire observation period using a linear regression model. A least-squares slope estimate of  $R_{atrophy}$  for each ROI from all available scans (for participants with  $\geq 4$  scans) was expressed as an annual rate of change. Conversion status was included in the model to ensure that our estimation of the anxiety effect was not attributable to greater  $R_{atrophy}$  due to AD.



## RESULTS

Sample characteristics and conversion status at study end are summarized in Table 1. Although all aMCI participants entered ADNI-1 with  $GDS \leq 5$  as per eligibility criteria, 8% of the sample reported  $GDS > 5$  on at least one study visit during the period of risk. The numbers of aMCI participants with any positive response for NPI-Q depression or anxiety during the period of risk were comparable (45% versus 41%). Similarly, NPI-Anxiety<sub>max</sub> and NPI-Dep<sub>max</sub> distributions were comparable.

In bivariate models, NPI-Anxiety<sub>max</sub>, NPI-Dep<sub>max</sub>, and GDS<sub>max</sub>, were significantly associated with rates of conversion to AD (Kaplan-Meier survival curves and HRs in figure), as were all cognitive variables (MMSE:HR=0.82,Wald  $z = -4.50$ ,  $p < 0.0001$ ; ADNI-MEM:HR=0.23,Wald  $z = -8.73$ ,  $p < 0.0001$ ; ADNI-EF: HR=0.52,Wald  $z = -6.42$ ,  $p < 0.0001$ ). The initial multivariable Cox proportional-hazards model included NPI-Anxiety<sub>max</sub>, NPI-Dep<sub>max</sub>, GDS<sub>max</sub>, MMSE, ADNI-MEM, ADNI-EF, age, sex, education, and Hachinski score. The final model after backward elimination included NPI-Anxiety<sub>max</sub>, ADNI-MEM, ADNI-EF, sex, and education and was confirmed with re-introduction of excluded variables. The HR<sub>p</sub> for anxiety (HR<sub>p-anxiety</sub>) in this model was 1.35 (summarized in Table 2). Neither NPI-Dep<sub>max</sub> nor GDS<sub>max</sub> met criteria for inclusion into the model but we report a second model with NPI-Dep<sub>max</sub> added in order to interpret the effects of anxiety on AD independent of depression (Table 2). Inclusion of NPI-Dep<sub>max</sub> resulted in reduction of the regression coefficient estimate for anxiety by 17%, and HR<sub>p-anxiety</sub> was minimally reduced. GDS<sub>max</sub> reduced the regression coefficient estimate for anxiety by less than 5%.

When either model was challenged by adding  $\Delta$ ADNI-MEM, the effect of anxiety on AD conversion remained significant and not substantively altered (HR<sub>p-anxiety</sub> remained at 1.30 and 1.27 (Wald  $z = 2.07$ ,  $p = .04$ ) for final or depression-adjusted model respectively).

### Neuroimaging predictors of AD risk

Baseline volumes of HC, AMYG, and EC were smaller in aMCI anxiety+ participants (Table 1). The multivariable Cox proportional-hazards model based on the subsample of 332 aMCI participants with MRI data was identical to the model indicated by the larger sample: retained predictors were NPI-Anxiety<sub>max</sub>, ADNI-MEM, ADNI-EF, sex, and education. Estimated HR<sub>p</sub> from this subsample was similar to the estimate from the full data set for each predictor. Estimated HR<sub>p-anxiety</sub> in this subsample was 1.25.

**Effect of baseline ROIs**—Table 3 summarizes the models with ICV and each baseline ROI added individually to the model. Baseline HC-vol, EC-vol, AMYG-vol, EC-vol, and EC-thickness, but not EC-surface, were strongly associated with AD risk. Although these neuroimaging associations reduced the magnitude of the effect of anxiety on AD conversion, anxiety remained a significant predictor of AD.

**Effect of change in ROIs over time**—Table 4 summarizes the models with  $\Delta$ ICV and each  $\Delta$ ROI added individually. Only  $\Delta$ EC-vol and  $\Delta$ EC-thickness were associated with AD. The regression coefficient estimate for anxiety was nominally increased with inclusion of  $\Delta$ ROIs in the model.

### Anxiety as a predictor of rate of atrophy within ROIs

$R_{\text{atrophy}}$  for each ROI in aMCI anxiety+ and aMCI anxiety- are summarized in Table 5. Anxiety status predicted  $R_{\text{atrophy}}$  of EC-vol. The association between anxiety and EC-thickness  $R_{\text{atrophy}}$  missed significance.

## DISCUSSION

Anxiety severity in aMCI increased rates of conversion to AD, independent of depression or extent of memory decline. The  $HR_p$  for anxiety was 1.33, indicating AD risk increased by 33%, 78%, and 135% for mild, moderate, and severe anxiety respectively. The association between anxiety and AD remained significant even with inclusion of baseline HC, AMYG, EC, or their extent of atrophy over time in the model. Further, anxiety status predicted greater annual rates of decrease in EC-vol.

Previous studies reported conflicting results on the contribution of depression to the association between anxiety and AD(19-21, 25) or did not document results(15-17). Further, the impact of depression developing after baseline on conversion has not been reported(15, 16). While depression is an exclusion criterion for enrollment into ADNI, 8% of aMCI participants developed clinically significant depression symptoms during follow-up, an incidence rate similar to the general population(50). Depression and anxiety-severity score distributions were comparable in the sample. Thus, it is noteworthy that the  $HR_p$  for anxiety remained significant and was minimally altered with depression severity included in the model ( $HR_p$  of 1.35 decreased to 1.33). Earlier studies reported similar HRs for baseline NPI-anxiety severity in MCI despite failing to adjust for depression(15, 18). A recent retrospective study in primary care demonstrated that anxiety attenuated the association between depression and dementia, and that AD risk was similar for anxiety comorbid with depression or anxiety alone(51). Collectively, these findings suggest that anxiety in aMCI increases risk of AD independently of depression.

If anxiety in aMCI accelerates progression towards AD, is it possible that anxiety is simply a prodromal symptom of AD - due to the presence of AD pathology in limbic brain regions(52) - or is anxiety a subjective reaction to worsening memory? Against these possibilities are the significant associations between anxiety and AD, over and beyond extent of memory decline, baseline ROI volumes, or ROI atrophy rate. We also observed no differences in frequency of endorsing the GDS question "Do you feel you have more problems with memory than most?" between aMCI anxiety+ and aMCI anxiety-, implying that anxiety was not simply related to concerns regarding memory.

Instead, we found that anxiety was associated with increased rate of atrophy within EC-vol, (an association with EC-thickness, specifically, missed significance), suggesting that anxiety may accelerate decline towards AD through effects on EC. Decreased EC-vol and EC-thickness were in turn associated with more rapid progression towards AD. The EC, along with perirhinal, parahippocampal cortices and HC, form the medial temporal lobe, which plays an integral role in memory formation(53). A link between anxiety and EC was previously demonstrated in a study of AD patients where NPI anxiety scores were inversely



correlated with metabolism in EC and anterior parahippocampal gyrus, after controlling for MMSE and NPI depression(54).

Contrary to hypotheses, anxiety was not associated with greater HC or AMYG atrophy rates. However, decreases in HC or AMYG-vols were also not predictive of conversion to AD. This pattern of results is consistent with pathological staging of AD, where neurofibrillary tangles appear initially in EC, followed by HC then AMYG in preclinical AD(52). Further, in healthy older adults, AD pathology as measured by amyloid- $\beta$  deposition was associated with abnormal neural activity during recall within EC, but not HC, suggesting that amyloid deposition is associated specifically with EC neuronal dysfunction(55).

One limitation to the current study is the use of NPI-Q in assessment of anxiety. Caregiver ratings may not accurately reflect the extent of anxiety experienced by individuals with aMCI. The NPI-Q also assesses a narrow range of anxiety symptoms, and does not characterize onset or chronicity. In a study focused on anxiety profiles in MCI, chronic severe anxiety, as opposed to recent onset, was associated with aMCI and correlated with memory impairment(56). Thus, it is possible that anxiety symptoms were underreported in the current study, resulting in a lower HR<sub>p</sub> estimate for anxiety and AD risk. Broader anxiety measures which are self- and clinician-rated are ideal, but challenging to implement in a large-scale, long-term observational study for which participant retention is critical. A second limitation relates to exclusion of comorbid depression on study entry in ADNI, a criterion implemented in other longitudinal AD studies since depression itself is associated with cognitive impairment(57). Thus, whether an association between anxiety and AD would be detected in aMCI with high levels of depression remains to be determined.

In summary, we demonstrated an association between anxiety symptoms in aMCI and rate of conversion to AD independent of depression, extent of memory loss, or atrophy of AD neuroimaging biomarkers. Further, anxiety in aMCI predicted greater EC atrophy rate, which itself was associated with AD risk. These findings are compatible with the hypothesis that anxiety is not a prodromal non-cognitive feature of AD, but may accelerate decline towards AD through direct or indirect effects on EC. Anxiety in aMCI has been understudied relative to other neuropsychiatric symptoms such as depression and apathy, despite well-established associations between stress, cortisol, and hippocampal integrity(30-32). This may, in part, be attributable to the tendency to subsume anxiety symptoms within the rubric of depression(58). Future research should clarify the clinical profile of anxiety and its association with AD in community samples of aMCI, and evaluate whether progression to AD can be delayed through routine monitoring and timely management of anxiety symptoms in aMCI; for example, using cognitive-behavioral interventions developed for anxiety in dementia(59).

## ACKNOWLEDGEMENTS

Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: Alzheimer's Association; Alzheimer's Drug Discovery Foundation; BioClinica, Inc.; Biogen Idec Inc.; Bristol-Myers Squibb Company; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; GE Healthcare; Innogenetics, N.V.; IXICO Ltd.; Janssen

Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Medpace, Inc.; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Synarc Inc.; and Takeda Pharmaceutical Company. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health ([www.fnih.org](http://www.fnih.org)). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Disease Cooperative Study at the University of California, San Diego. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of California, Los Angeles. This research was also supported by NIH grants P30 AG010129 and K01 AG030514.

LM has received funding through the Geoffrey H. Wood Foundation, Baycrest Department of Psychiatry Academic Development Fund, and the Ministry of Health and Long-Term Care Academic Funding Plan Innovation Fund.


## REFERENCES

1. Albert MS, DeKosky ST, Dickson D, et al. The diagnosis of mild cognitive impairment due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement*. 2011; 7:270–279. [PubMed: 21514249]
2. Petersen RC, Doody R, Kurz A, et al. Current concepts in mild cognitive impairment. *Archives of neurology*. 2001; 58:1985–1992. [PubMed: 11735772]
3. Petersen RC, Smith GE, Waring SC, et al. Mild cognitive impairment: clinical characterization and outcome. *Archives of neurology*. 1999; 56:303–308. [PubMed: 10190820]
4. Ganguli M, Snitz BE, Saxton JA, et al. Outcomes of mild cognitive impairment by definition: a population study. *Archives of neurology*. 2011; 68:761–767. [PubMed: 21670400]
5. Cui Y, Liu B, Luo S, et al. Identification of conversion from mild cognitive impairment to Alzheimer's disease using multivariate predictors. *PLoS One*. 2011; 6:e21896. [PubMed: 21814561]
6. Davatzikos C, Bhatt P, Shaw LM, et al. Prediction of MCI to AD conversion, via MRI, CSF biomarkers, and pattern classification. *Neurobiol Aging*. 2011; 32:2322, e2319–2327. [PubMed: 20594615]
7. Ewers M, Walsh C, Trojanowski JQ, et al. Prediction of conversion from mild cognitive impairment to Alzheimer's disease dementia based upon biomarkers and neuropsychological test performance. *Neurobiol Aging*. 2012; 33:1203–1214. [PubMed: 21159408]
8. Weiner MW, Veitch DP, Aisen PS, et al. The Alzheimer's Disease Neuroimaging Initiative: a review of papers published since its inception. *Alzheimers Dement*. 2012; 8:S1–68. [PubMed: 22047634]
9. Apostolova LG, Cummings JL. Neuropsychiatric manifestations in mild cognitive impairment: a systematic review of the literature. *Dement Geriatr Cogn Disord*. 2008; 25:115–126. [PubMed: 18087152]
10. Hwang TJ, Masterman DL, Ortiz F, et al. Mild cognitive impairment is associated with characteristic neuropsychiatric symptoms. *Alzheimer Dis Assoc Disord*. 2004; 18:17–21. [PubMed: 15195459]
11. Lyketsos CG, Lopez O, Jones B, et al. Prevalence of neuropsychiatric symptoms in dementia and mild cognitive impairment: results from the cardiovascular health study. *JAMA*. 2002; 288:1475–1483. [PubMed: 12243634]
12. Steffens DC. Separating mood disturbance from mild cognitive impairment in geriatric depression. *Int Rev Psychiatry*. 2008; 20:374–381. [PubMed: 18925486]
13. Peters ME, Rosenberg PB, Steinberg M, et al. Prevalence of neuropsychiatric symptoms in CIND and its subtypes: the Cache County Study. *The American journal of geriatric psychiatry : official journal of the American Association for Geriatric Psychiatry*. 2012; 20:416–424. [PubMed: 22522960]
14. Ownby RL, Crocco E, Acevedo A, et al. Depression and risk for Alzheimer disease: systematic review, meta-analysis, and meta-regression analysis. *Archives of general psychiatry*. 2006; 63:530–538. [PubMed: 16651510]

15. Wadsworth LP, Lorus N, Donovan NJ, et al. Neuropsychiatric symptoms and global functional impairment along the Alzheimer's continuum. *Dement Geriatr Cogn Disord*. 2012; 34:96–111. [PubMed: 22922821]
16. Somme J, Fernandez-Martinez M, Molano A, et al. Neuropsychiatric symptoms in amnesic mild cognitive impairment: increased risk and faster progression to dementia. *Curr Alzheimer Res*. 2013; 10:86–94. [PubMed: 23016837]
17. Palmer K, Berger AK, Monastero R, et al. Predictors of progression from mild cognitive impairment to Alzheimer disease. *Neurology*. 2007; 68:1596–1602. [PubMed: 17485646]
18. Rosenberg PB, Mielke MM, Appleby BS, et al. The association of neuropsychiatric symptoms in MCI with incident dementia and Alzheimer disease. *The American journal of geriatric psychiatry : official journal of the American Association for Geriatric Psychiatry*. 2013; 21:685–695. [PubMed: 23567400]
19. Liu HC, Wang PN, Wang HC, et al. Conversion to dementia from questionable dementia in an ethnic Chinese population. *J Geriatr Psychiatry Neurol*. 2007; 20:76–83. [PubMed: 17548776]
20. Devier DJ, Pelton GH, Tabert MH, et al. The impact of anxiety on conversion from mild cognitive impairment to Alzheimer's disease. *International journal of geriatric psychiatry*. 2009; 24:1335–1342. [PubMed: 19319929]
21. Ramakers IH, Visser PJ, Aalten P, et al. Affective symptoms as predictors of Alzheimer's disease in subjects with mild cognitive impairment: a 10-year follow-up study. *Psychol Med*. 2010; 40:1193–1201. [PubMed: 19903364]
22. Brodaty H, Heffernan M, Draper B, et al. Neuropsychiatric symptoms in older people with and without cognitive impairment. *J Alzheimers Dis*. 2012; 31:411–420. [PubMed: 22571979]
23. Gallagher D, Coen R, Kilroy D, et al. Anxiety and behavioural disturbance as markers of prodromal Alzheimer's disease in patients with mild cognitive impairment. *International journal of geriatric psychiatry*. 2011; 26:166–172. [PubMed: 21229599]
24. Chan WC, Lam LC, Tam CW, et al. Neuropsychiatric symptoms are associated with increased risks of progression to dementia: a 2-year prospective study of 321 Chinese older persons with mild cognitive impairment. *Age Ageing*. 2011; 40:30–35. [PubMed: 21106558]
25. Robert PH, Berr C, Volteau M, et al. Apathy in patients with mild cognitive impairment and the risk of developing dementia of Alzheimer's disease: a one-year follow-up study. *Clin Neurol Neurosurg*. 2006; 108:733–736. [PubMed: 16567037]
26. Kessler RC, Chiu WT, Demler O, et al. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of general psychiatry*. 2005; 62:617–627. [PubMed: 15939839]
27. Kessler RC, Gruber M, Hettema JM, et al. Co-morbid major depression and generalized anxiety disorders in the National Comorbidity Survey follow-up. *Psychol Med*. 2008; 38:365–374. [PubMed: 18047766]
28. Ramakers IH, Verhey FR, Scheltens P, et al. Anxiety is related to Alzheimer cerebrospinal fluid markers in subjects with mild cognitive impairment. *Psychol Med*. 2013; 43:911–920. [PubMed: 22954311]
29. Green KN, Billings LM, Roozendaal B, et al. Glucocorticoids increase amyloid-beta and tau pathology in a mouse model of Alzheimer's disease. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2006; 26:9047–9056. [PubMed: 16943563]
30. Lupien SJ, de Leon M, de Santi S, et al. Cortisol levels during human aging predict hippocampal atrophy and memory deficits. *Nat Neurosci*. 1998; 1:69–73. [PubMed: 10195112]
31. Sapolsky RM, Uno H, Rebert CS, et al. Hippocampal damage associated with prolonged glucocorticoid exposure in primates. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 1990; 10:2897–2902. [PubMed: 2398367]
32. Erickson K, Drevets W, Schulkin J. Glucocorticoid regulation of diverse cognitive functions in normal and pathological emotional states. *Neurosci Biobehav Rev*. 2003; 27:233–246. [PubMed: 12788335]
33. Mantella RC, Butters MA, Amico JA, et al. Salivary cortisol is associated with diagnosis and severity of late-life generalized anxiety disorder. *Psychoneuroendocrinology*. 2008; 33:773–781. [PubMed: 18407426]

34. McEwen BS, Stellar E. Stress and the individual. Mechanisms leading to disease. *Archives of internal medicine*. 1993; 153:2093–2101. [PubMed: 8379800]
35. Augustinack JC, Huber KE, Postelnicu GM, et al. Entorhinal verrucae geometry is coincident and correlates with Alzheimer's lesions: a combined neuropathology and high-resolution ex vivo MRI analysis. *Acta neuropathologica*. 2012; 123:85–96. [PubMed: 22160360]
36. Mueller SG, Weiner MW, Thal LJ, et al. The Alzheimer's disease neuroimaging initiative. *Neuroimaging Clin N Am*. 2005; 15:869–877. xi–xii. [PubMed: 16443497]
37. Yesavage JA, Brink TL, Rose TL, et al. Development and validation of a geriatric depression screening scale: a preliminary report. *J Psychiatr Res*. 1982; 17:37–49. [PubMed: 7183759]
38. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975; 12:189–198. [PubMed: 1202204]
39. Berg L. Clinical Dementia Rating (CDR). *Psychopharmacol Bull*. 1988; 24:637–639. [PubMed: 3249765]
40. Kaufer DI, Cummings JL, Ketchel P, et al. Validation of the NPI-Q, a brief clinical form of the Neuropsychiatric Inventory. *The Journal of neuropsychiatry and clinical neurosciences*. 2000; 12:233–239. [PubMed: 11001602]
41. Rosen WG, Terry RD, Fuld PA, et al. Pathological verification of ischemic score in differentiation of dementias. *Annals of neurology*. 1980; 7:486–488. [PubMed: 7396427]
42. Crane PK, Carle A, Gibbons LE, et al. Development and assessment of a composite score for memory in the Alzheimer's Disease Neuroimaging Initiative (ADNI). *Brain imaging and behavior*. 2012; 6:502–516. [PubMed: 22782295]
43. Rey, A. Psychological appraisal of children with cerebral deficits. Harvard University Press; Cambridge, Massachusetts: 1964. Auditory Verbal Learning Test.. 196 I
44. Mohs RC, Knopman D, Petersen RC, et al. Development of cognitive instruments for use in clinical trials of antidementia drugs: additions to the Alzheimer's Disease Assessment Scale that broaden its scope. *The Alzheimer's Disease Cooperative Study*. *Alzheimer Dis Assoc Disord*. 1997; 11(Suppl 2):S13–21. [PubMed: 9236948]
45. Wechsler, D. WMS-R: Wechsler Memory Scale-Revised. Psychological Corporation; 1987.
46. Gibbons LE, Carle AC, Mackin RS, et al. A composite score for executive functioning, validated in Alzheimer's Disease Neuroimaging Initiative (ADNI) participants with baseline mild cognitive impairment. *Brain imaging and behavior*. 2012; 6:517–527. [PubMed: 22644789]
47. Reuter M, Rosas HD, Fischl B. Highly accurate inverse consistent registration: a robust approach. *Neuroimage*. 2010; 53:1181–1196. [PubMed: 20637289]
48. Reuter M, Schmansky NJ, Rosas HD, et al. Within-subject template estimation for unbiased longitudinal image analysis. *Neuroimage*. 2012; 61:1402–1418. [PubMed: 22430496]
49. Fisher LD, Lin DY. Time-dependent covariates in the Cox proportional-hazards regression model. *Annual review of public health*. 1999; 20:145–157.
50. Buchtemann D, Lupp M, Bramesfeld A, et al. Incidence of late-life depression: a systematic review. *Journal of affective disorders*. 2012; 142:172–179. [PubMed: 22940498]
51. Burton C, Campbell P, Jordan K, et al. The association of anxiety and depression with future dementia diagnosis: a case-control study in primary care. *Fam Pract*. 2013; 30:25–30. [PubMed: 22915794]
52. Braak H, Braak E. Neuropathological staging of Alzheimer-related changes. *Acta neuropathologica*. 1991; 82:239–259. [PubMed: 1759558]
53. Squire LR, Stark CE, Clark RE. The medial temporal lobe. *Annual review of neuroscience*. 2004; 27:279–306.
54. Hashimoto H, Monserratt L, Nguyen P, et al. Anxiety and regional cortical glucose metabolism in patients with Alzheimer's disease. *The Journal of neuropsychiatry and clinical neurosciences*. 2006; 18:521–528. [PubMed: 17135378]
55. Huijbers W, Mormino EC, Wigman SE, et al. Amyloid deposition is linked to aberrant entorhinal activity among cognitively normal older adults. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2014; 34:5200–5210. [PubMed: 24719099]

56. Andreescu C, Teverovsky E, Fu B, et al. Old worries and new anxieties: behavioral symptoms and mild cognitive impairment in a population study. *The American journal of geriatric psychiatry : official journal of the American Association for Geriatric Psychiatry*. 2014; 22:274–284. [PubMed: 23759435]
57. Steffens DC, Otey E, Alexopoulos GS, et al. Perspectives on depression, mild cognitive impairment, and cognitive decline. *Archives of general psychiatry*. 2006; 63:130–138. [PubMed: 16461855]
58. Roth M, Gurney C, Garside RF, et al. Studies in the classification of affective disorders. The relationship between anxiety states and depressive illnesses. I. *The British journal of psychiatry : the journal of mental science*. 1972; 121:147–161. [PubMed: 5072240]
59. Stanley MA, Calleo J, Bush AL, et al. The peaceful mind program: a pilot test of a cognitive-behavioral therapy-based intervention for anxious patients with dementia. *The American journal of geriatric psychiatry : official journal of the American Association for Geriatric Psychiatry*. 2013; 21:696–708. [PubMed: 23567399]

**FIGURE.**

Kaplan-Meier survival curves of progression from aMCI to AD dementia as predicted by NPI-Q anxiety severity (left graph), NPI-Q depression severity (middle graph), and GDS (right graph). Survival indicates absence of progression from aMCI to AD dementia.



Table 1

Demographic, clinical, and neuroimaging characteristics of aMCI participants

Variable	Total MCI sample	MCI without anxiety	MCI with anxiety	$ d $ <sup>*</sup>
<b>Number</b>	376	220	156	
Age at baseline, years	75.0 (7.26)	75.2 (7.55)	74.7 (6.84)	0.07
Male, %	64	70	56	0.30
Education, years	15.6 (3.03)	15.6 (3.08)	15.6 (2.98)	<0.01
Modified Hachinski ischemia score	0.63 (0.71)	0.64 (0.69)	0.63 (0.76)	0.01
Duration of follow-up/period of observation, years	31.0 (11.4)	31.2 (11.7)	30.6 (11.0)	0.06
% followed for minimum of 24 months	80.3	80	80.7	
Number converted to AD	165	87	78	
<b>Cognition</b>				
MMSE at study entry	27.0 (1.77)	27.0 (1.73)	26.9 (1.84)	0.06
Memory composite score (ADNI-MEM) at baseline	-0.0997 (0.57)	-0.0766 (0.57)	-0.1322 (0.58)	0.10
Executive Function composite score (ADNI-EF) at baseline	-0.0581 (0.76)	-0.0578 (0.80)	-0.0584 (0.70)	<0.01
<b>Anxiety</b>				
Baseline NPI-Q anxiety response (yes/no), % Yes	18	0	44	
%mild/moderate/severe			68/29/3	
<b>Depression</b>				
Baseline GDS score, possible range 0-15	1.53 (1.36)	1.38 (1.29)	1.74 (1.42)	0.27
Highest GDS score during period of observation, mean (sd, range)	2.59 (2.25)	2.24 (1.95)	3.08 (2.54)	0.38
Highest GDS score during period of observation, % participants	7.9	6.8	9.6	
Baseline NPI-Q depression response (yes/no), % Yes	19	14	26	0.31
Any positive NPI-Q depression response during period of observation, % Yes	45	33	62	0.61
Highest NPI-Q depression severity score during period of observation, %mild/moderate/severe	65/32/3	78/22/0	55/40/5	
<b>Baseline values for AD neuroimaging biomarkers</b>				
<b>Number</b>	192		140	
ICV ( $\times 10^3$ mL)	1 586 (174)		1 556 (158)	0.18
Amygdala vol (mL)	1 027 (202)		950 (202)	0.38
Hippocampal vol (mL)	2 967 (518)		2 827 (459)	0.28
EC vol (mL)	1 735 (409)		1 612 (358)	0.32

Variable	Total MCI sample	MCI without anxiety	MCI with anxiety	*  d
EC surface area (mm <sup>2</sup> )		359 (78)	370 (78)	0.14
EC thickness (mm)		3.14 (0.521)	3.11 (0.520)	0.06

Summary statistics are mean (sd) for continuous measures and % for categorical measures

\* Cohen's *d* statistic to quantify difference between aMCI anxiety+ and noaMCI anxiety-. small (*d* = 0.20), medium (*d* = 0.50), large (*d* = 0.80).

Author Manuscript

*Am J Geriatr Psychiatry*. Author manuscript; available in PMC 2016 May 01.

**Table 3**  
Effect of individual baseline structural neuroimaging measures on hazard for AD and on hazard ratio ( $HR_p$ ) for anxiety.

Region of interest (ROI)	$HR_p$ for ROI	Wald z-value	p-value	$HR_p$ for Anxiety with inclusion of ROI in the model <sup><math>\delta</math></sup>
ICV (mL) <sup>*</sup>	1.00	+0.71	0.48	1.25
HC volume (mL)	0.52	-2.88	0.004	1.20
Amygdala volume (mL)	0.18	-3.12	0.002	1.18
EC volume (mL)	0.42	-3.35	0.0008	1.16
EC thickness (mm)	0.64	-2.55	0.01	1.21
EC surface area (cm <sup>2</sup> )	0.94	-0.54	0.59	1.24

<sup>\*</sup> included in all models

<sup>$\delta$</sup>  Cox proportional-hazards models (130 events, n=332) additionally included sex, education, baseline memory, baseline executive function (see text).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 4**

Effect of change in ROI ( $\Delta$ ROI)<sup>†</sup> at each visit from baseline on hazard of AD and on hazard ratio (HR<sub>p</sub>) for anxiety. (ICV = intracranial volume, HC = hippocampus, EC = entorhinal cortex)

Region of interest (ROI)	HR <sub>p</sub>	Wald z-value	p-value	HR <sub>p</sub> for Anxiety with inclusion of $\Delta$ ROI in the model <sup>δ</sup>
$\Delta$ ICV (mL) <sup>*</sup>	1.01	+1.57	0.12	1.27
$\Delta$ HC (mL)	0.33	-1.09	0.28	1.27
$\Delta$ Amygdala (mL)	0.27	-0.98	0.33	1.28
$\Delta$ EC volume (mL)	0.08	-2.23	0.03	1.29
$\Delta$ EC thickness (mm)	0.12	-3.28	0.001	1.29
$\Delta$ EC surface area (cm <sup>2</sup> )	0.74	-0.78	0.43	1.27

<sup>†</sup> see text for calculation of  $\Delta$ ROI

<sup>\*</sup> included in all models

<sup>δ</sup> Cox proportional-hazards models (130 events, n=332) additionally included sex, education, baseline memory, baseline executive function (see text).

Mean (and standard deviation) for rate of change ( $R_{atrophy}$ ) in AD structural neuroimaging biomarkers by anxiety status in aMCI. (ICV = intracranial volume, HC = hippocampus, EC = entorhinal cortex)

$R_{atrophy}$ within region of interest	MCI anxiety-	MCI anxiety+	t-statistic <sup>†</sup>
Number	132	104	
ICV	+1.78 (sd = 7.43)	+2.34 (sd = 6.69)	+0.55 (p = 0.58)
HC volume	-0.072 (sd = 0.054)	-0.086 (sd = 0.053)	-1.36 (p = 0.18)
Amygdala volume	-0.023 (sd = 0.035)	-0.022 (sd = 0.035)	+0.35 (p = 0.73)
EC volume	-0.020 (sd = 0.047)	-0.070 (sd = 0.049)	-2.55 (p = 0.01)
EC thickness	-0.078 (sd = 0.080)	-0.106 (sd = 0.084)	-1.93 (p = 0.0546)
EC surface area	-0.035 (sd = 0.097)	-0.047 (sd = 0.104)	-0.68 (p = 0.50)

<sup>†</sup> conversion status included as a covariate, df = 233